arXiv:1206.2226v1 [math.GT] 11 Jun 2012

On stable Khovanov homology of torus knots

E. Gorsky, A. Oblomkov, J. Rasmussen

Abstract

We conjecture that the stable Khovanov homology of torus knots can be described Koszul homology of an explicit non-regular sequence of quadratic polynomials. The corresponding Poincaré series turns out to be related to the Rogers-Ramanujan identity.

1 Introduction

In [23] M. Khovanov constructed a knot homology theory which categorifies the Jones polynomial using a combinatorial construction in terms of a knot projection. Following the early computations of Bar-Natan and Shumakovitch [3], [36], [6], it became evident that the torus knots $T(n,m)$ had "interesting" Khovanov homology, in the sense that their homology was much larger than might have been guessed from looking at the corresponding Jones polynomial, had torsion of high order, *etc.* Further advances in computation, most notably Bar-Natan's geometric Khovanov homology [5], have enabled us to calculate Khovanov homology of torus knots up through $T(7,n)$, where n is relatively large [6], [37]. These calculations have tended to confirm our first impression of overall complexity.

Nevertheless, there are indications that the Khovanov homology of torus knots is not only interesting, but may be important to our understanding of Khovanov homology as a whole. The first result in this direction is the theorem of Stošić [39], who proved that if we fix n and allow m to vary, then (after a suitable renormalization), the groups $\mathrm{Kh}(T(n,m))$ tend to a well-defined limit, which we denote by $\mathrm{Kh}(T(n,\infty))$. More recently, Rozansky [35] has shown that the Khovanov complex of the infinite torus braid provides a categorified version of the Jones-Wenzl projector, and thus should play an important role in the theory of colored Khovanov homology [15], [18] [44]. In this framework, $\mathrm{Kh}(n,\infty)$ appears as the n-colored Khovanov homology of the unknot.

In this paper, we propose a conjectural description of $\mathrm{Kh}(T(n,\infty))$ for all n:

Conjecture 1 *The unreduced stable Khovanov homology* $\mathrm{Kh}(T(n,\infty))$ *is dual to the homology of the differential graded algebra generated by even variables* $x_0,\ldots,x_{n-1}$ *and odd variables* $\xi_0,\ldots,\xi_{n-1}$, *equipped with the differential* d_2 *defined by*

$$d_2(\xi_k)=\sum_{i=0}^{k}x_ix_{k-i} \quad \textit{and} \quad d_2(x_k)=0.$$

Equivalently, this is the Koszul complex determined by the (nonregular) sequence $d_2(\xi_k)$ *for* $k=0,\ldots,n-1$.

Remark 2 *The homology of the chain complex described in the conjecture should be the $\mathfrak{sl}(2)$ Khovanov-Rozansky homology [26] of $T(n,\infty)$. This, in turn, is dual to the ordinary Khovanov homology (in the usual sense that they are homologies of dual chain complexes.)*

Khovanov homology is bigraded; it is equipped with both *polynomial* (q) and *homological* (t) gradings. With respect to the usual normalizations for these gradings, the generators x_k and ξ_k are graded as follows:

$$\deg(x_k) = q^{2k+2}t^{2k}, \quad \deg(\xi_k) = q^{2k+4}t^{2k+1}.$$

The differential d_2 preserves q-grading and lowers the t-grading by 1.

The conjecture was motivated by our work [17] on the relation between the HOMFLY-PT [27] homology of torus knots and the representation theory of the rational Cherednik algebra. More specifically, it is known [9], [33] that one can pass from the HOMFLY-PT homology of a knot K to its $\mathfrak{sl}(N)$ Khovanov-Rozansky homology [26] by means of a spectral sequence. The main conjecture of [17] relates the HOMFLY-PT homology of $T(n,m)$ to the representation theory of the rational Cherednik algebra. On the right-hand side of this equivalence, it is possible to construct certain natural differentials which we believe should correspond to the differentials needed to pass to the $\mathfrak{sl}(N)$ homology. We arrived at the conjecture above by computing these representation-theoretic differentials for $N = 2$ in the limiting case $m \to \infty$.

We hope that the conjecture will be interesting purely in the context of Khovanov homology, and will adopt this perspective for the remainder of this paper.

Definition 3 *We will denote the homology of d_2 by $\mathrm{Kh}_{alg}(n,\infty)$.*

An immediate consequence of the conjecture is that $\mathrm{Kh}_{alg}(n,\infty)$ is especially simple if we use $\mathbb{Z}/2$ coefficients:

Theorem 4 $\mathrm{Kh}_{alg}(n,\infty;\mathbb{Z}/2)$ *has the following Poincaré series:*

$$P_n(q,t;\mathbb{Z}_2) = \prod_{i=0}^{n-1} \frac{(1+q^{2i+4}t^{2i+1})}{(1-q^{2i+2}t^{2i})} \prod_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \frac{(1-q^{4i+4}t^{4i})}{(1+q^{4i+4}t^{4i+1})}.$$

With rational coefficients, the homology is more complicated. In section 3.3, we construct some explicit elements of $\mathrm{Kh}_{alg}(n,\infty;\mathbb{Q})$, as well as some relations which they satisfy. This leads to the following

Conjecture 5 *As an algebra over $\mathbb{Q}$, $\mathrm{Kh}_{alg}(n,\infty)$ has a presentation with generators $x_0,\dots,x_{n-1}$ and $\mu_0,\dots,\mu_{n-2}$ (where μ_i has bidegree $q^{2i+8}t^{2i+3}$) and relations of the form*

$$x(z)^2 = 0, \quad x(z)\mu(z) = 0, \quad \ddot{x}(z)\mu(z) - \dot{x}(z)\dot{\mu}(z) = 0, \quad \mu(z)\dot{\mu}(z) = 0 \qquad (1)$$

where $x(z) = \sum_{i=0}^{n-1} x_i z^i$ $\mu(z) = \sum_{i=0}^{n-2} \mu_i z^i$, and each equation above is to be interpreted modulo z^n.

Following the ideas of B. Feigin and A. Stoyanovsky ([13], see also [14], [30]), we derive a conjectural explicit formula for the Poincaré polynomial of $\mathrm{Kh}(T(n,\infty))$. Feigin and Stoyanovsky studied the structure of the coinvariants for the integrable representation of $\widehat{sl}(2)$ at level 1 using the vertex operator equations of Lepowsky and Primc ([29]), which turn out to be analogous to our Koszul differential. The resulting stable homology are described by the equation (2) for the unreduced theory, and by the equation (19) for the reduced theory. They generalize the result of Feigin and Stoyanovsky, which is itself a generalization of the Rogers-Ramanujan identity ([2]).

Conjecture 6 *The Poincaré series of* $\mathrm{Kh}_{alg}(n,\infty;\mathbb{Q})$ *can be expressed as*

$$P_n(q,t) = \frac{1}{\prod_{k=1}^{n}(1-q^{2k}t^{2k-2})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}(1-q^{2k}t^{2k-2})\times \tag{2}$$

$$\times\prod_{k=3p+1}^{n-1}(1+q^{2k+6}t^{2k+1})\prod_{k=1}^{2p-1}(1+q^{2k+2}t^{2k-1})\times$$

$$[q^{5p^2+p}t^{5p^2-3p}(1+\chi_p^+q^{6p+4}t^{6p-1})(1+q^{6p+6}t^{6p+1})\binom{n-2p+1}{p}_z+$$

$$q^{5p^2+7p+2}t^{5p^2+3p-1}(1+q^{6p+6}t^{6p+1})(1-q^{2p+2}t^{2p})\binom{n-2p}{p}_z-$$

$$q^{5p^2+9p+4}t^{5p^2+5p}(1+q^{2p+2}t^{2p+1})(1+\chi_p^+q^{4p+2}t^{4p-1})\binom{n-2p-1}{p}_z],$$

where $\chi_p^+=0$ *when* $p=0$, $\chi_p^+=1$ *for* $p>0$,

$$z=q^2t^2, \quad \textit{and} \quad \binom{a}{b}_z=\frac{(1-z)\cdots(1-z^a)}{(1-z)\cdots(1-z^b)(1-z)\cdots(1-z^{a-b})}.$$

The following conjecture is due to A. Shumakovitch and P. Turner:

Conjecture 7 *([37]) Let* $\mathcal{K}_n(q,t)$ *denote the Poincaré polynomials for the Khovanov homology for the* $(n,n+1)$ *torus knot. Then*

$$\mathcal{K}_n(q,t)=\mathcal{K}_{n-1}(q,t)+\mathcal{K}_{n-2}(q,t)q^{2n}t^{2n-2}+\mathcal{K}_{n-3}(q,t)q^{2n+4}t^{2n-1}. \tag{3}$$

We prove the following

Theorem 8 *If* $K_n(q,t)$ *is given by the recursion relation (3) with the appropriate initial conditions and* $P_n(q,t)$ *is the Hilbert series of the algebra described in Conjecture 5, then*

$$\lim_{n\to\infty}K_n(q,t)=\lim_{n\to\infty}P_n(q,t).$$

Finally, we describe an intriguing connection to the physical models of coloured homology proposed by S. Gukov, J. Walcher and M. Stošić ([18],[19]). In these models the homology of the unknot is constructed as the Milnor algebra of the certain potential W_{phys} with an isolated singularity.

Theorem 9 *The homology of d_2 is isomorphic to the Hochschild homology of the category of matrix factorizations of a certain potential W. The potential W has a non-isolated singularity (for $n > 1$) and coincided with a bihomogeneous part of W_{phys} of bidegree $(2n+4, 2n-2)$.*

We are grateful to B. Feigin, S. Gukov, M. Hagencamp, M. Khovanov, A. Kirillov Jr., S. Loktev, L. Rozansky, M. Stošić, J. Sussan, O. Viro, and V. Shende for the useful discussions. Special thanks to A. Shumakovitch for providing us with the valuable Khovanov homology data and explaining the Conjecture 7. Most of the computations of the Koszul homology were done using `Singular`, a computer algebra system ([8]). The research of E. G. was partially supported by the grants RFBR-10-01-00678, NSh-8462.2010.1 and the Simons foundation.

2 Evidence for the Conjecture

In this section, we outline the evidence in support of Conjecture 1. We verify that the conjecture holds for $T(n,\infty)$ in the cases $n = 2, 3$, where the Khovanov homology is well-understood. We then discuss the computational evidence for larger values of n.

We define the stable Khovanov homology by

$$\mathrm{Kh}(T(n,\infty)) := \lim_{m\to\infty} q^{-(n-1)(m-1)+1}\,\mathrm{Kh}(T(n,m)).$$

It is a theorem of Stošić [39] that this limit exists. The stable homology is normalized so that its Poincaré polynomial is a polynomial in q and t (rather than just a Laurent polynomial), with constant term 1.

2.1 $T(2,\infty)$

The Khovanov homology of $T(2,\infty)$ is well-known. In the language of [4], it can be viewed as dual to the homology of the following chain complex:



This picture has the following meaning. The Khovanov homology of the unknot is two-dimensional; as an algebra it can be described as $H_0 = \mathbb{C}[x_0]/(x_0^2)$. This algebra carries a comultiplication μ defined by the equations:

$$\mu : H_0 \to H_0 \otimes H_0, \quad \mu(1) = 1 \otimes x_0 + x_0 \otimes 1, \quad \mu(x_0) = 0.$$

Recall that the q-degree of x_0 equals to 2.

The complex is generated by the infinite number of copies of H_0 in t-degrees $0, 2, 3, 4, \ldots$. The q-grading in a k-th copy is shifted by $2k$. The maps between $2k+1$-th copy and $2k$-th are given by the cobordism on the picture, which can be presented as a composition of the comultiplication and multiplication:

$$H_0 \longleftarrow 0 \longleftarrow H_0[2]\{4\} \xleftarrow{m\circ\mu} H_0[3]\{6\} \xleftarrow{0} H_0[4]\{8\} \xleftarrow{m\circ\mu} H_0[5]\{10\} \xleftarrow{0} \cdots \tag{4}$$

Remark that $m \circ \mu$ coincided with the multiplication by $2x_0$, and let us introduce two formal variables x_1 of bidegree q^4t^2 and ξ_1 of bidegree q^6t^3 assuming ξ_1 to be odd. In other words, we identify $H_0[2k]\{4k\}$ with $x_1^k \cdot H_0$ and $H_0[2k+3]\{4k+6\}$ with $x_1^k\xi_1 \cdot H_0$

Then the complex (4) can be rewritten as an algebra $H_0[x_1, \xi_1]$ with the differential $d(\xi_1) = 2x_0x_1$, which is equivalent to our Koszul model.

2.2 $T(3, \infty)$

With rational coefficients, the Khovanov homology of $T(3, n)$ was computed by Turner [42]. The Poincaré polynomial of the stable homology is

$$P_{3,\infty} = (1 + q^2 + q^4t^2) + q^6t^3 \left(\frac{q^2 + t + q^2t + q^4t^2 + q^6t^2 + q^4t^3}{1 - q^6t^4} \right).$$

$\mathrm{Kh}_{alg}(3, \infty; \mathbb{Q})$ is computed in section 3.2. Its Poincaré polynomial is easily seen to agree with the one given above.

2.3 $\mathbb{Z}_2$ coefficients

In many cases, the Khovanov homology with $\mathbb{Z}_2$ coefficients is simpler that the homology with rational coefficients. It turns out that the stable answers become especially simple if we work over $\mathbb{Z}_2$.

Theorem 10 $\mathrm{Kh}_{alg}\,(n, \infty; \mathbb{Z}/2)$ *has the following Poincaré series:*

$$P_n(q, t; \mathbb{Z}_2) = \prod_{i=0}^{n-1} \frac{(1 + q^{2i+4}t^{2i+1})}{(1 - q^{2i+2}t^{2i})} \prod_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \frac{(1 - q^{4i+4}t^{4i})}{(1 + q^{4i+4}t^{4i+1})}.$$

Proof. In characteristic 2 the differential d_2 degenerates to the following form:

$$d_2(\xi_{2k}) = x_k^2, \quad d_2(\xi_{2k+1}) = 0.$$

Therefore for every $0 \le k \le \lfloor \frac{n-1}{2} \rfloor$ the odd generator ξ_{2i} kills x_i^2 in the homology. ■

We used JavaKh [6] to verify that $\mathrm{Kh}_{alg}(n, \infty; \mathbb{Z}/2)$ agrees with $\mathrm{Kh}(T(n, m))$ in the stable range (q-degree $\le 2m$) for $(n, m) = (3, 50), (4, 49), (5, 29)$.

2.4 $\mathbb{Q}$ coefficients

If we use rational coefficients, the structure of Kh_{alg} is more complicated (*c.f.* the conjectures in Section 3 below.) The rational Khovanov homology of torus knots has been extensively computed by Shumakovitch [37]. By comparing with his results, we have verified Conjecture 1 in the stable range up to $(n, m) = (7, 20)$.

In testing Conjecture 1, it is important to check that the predictions it makes about Khovanov homology can be distinguished from the ones we would get if we replaced the differential $d_2(\xi_k) = \sum x_i x_{k-i}$ with $d_2'(\xi_k) = \sum \alpha_{ik} x_i x_{k-i}$ for generic values of α_{ik}. In addition to the information on torsion discussed in this section, we can see evidence of this fact with rational coefficients in the case $n = 7$. As discussed in Remark 22 below, for generic α_{ik}, the homology with respect to d_2' has smaller dimension that the homology with respect to d_2, and the latter groups agree with the actual Khovanov homology.

More precisely, Remark 22 shows that for our choice of α_{ik} the homology in bidegree $q^{18}t^{13}$ is one-dimensional and spanned by the generator μ_5, while it vanishes for a generic choice. Theorem 6 in [40] states that the (homological) degree i homology of (p, q) torus knot are isomorphic to the ones of $(p, q + 1)$ torus knot, if $i < p + q - 2$. Since $13 < 7 + 9 - 2$, we conclude that the coefficient at t^{13} in stable homology of $(7, \infty)$ torus knot coincides with the same coefficient for $(7, 9)$ knot. The Poincaré polynomial for the Khovanov homology of the latter knot is presented in the Appendix C, and the term $q^{18}t^{13}$ is present.

2.5 $\mathbb{Z}_p$ torsion

The odd torsion in Khovanov homology was studied in [1] and [38] for some classes of knots, and [5] shows how complicated the torsion can be on the example of $(7, 8)$ knot. It was suggested that Khovanov homology can have torsion of arbitrarily large order. The following calculation provides support for this claim, as well as some additional evidence in favor of Conjecture 1.

Theorem 11 *Let $p > 3$ be a prime number. Then $\mathrm{Kh}_{alg}(p, \infty)$ has nontrivial $\mathbb{Z}_p$-torsion at bidegree $q^{2p+6}t^{2p}$.*

Proof. Consider an element

$$m = \sum_{i=1}^{p-1} (3i - p) x_i \xi_{p-i} = \sum_{i+j=p, 1<i<p-1} (2i - j) x_i \xi_j,$$

then

$$d_2(m) = \sum_{i+j+k=p, 1<i<p-1} (2i-j-k) x_i x_j x_k = \sum_{i+j+k=p} (2i-j-k) x_i x_j x_k - 2p x_p x_0^2 - p x_0 \sum_{j+k=p} x_j x_k.$$

Since the first sum vanishes, we have

$$d_2(m) = -2p x_p x_0^2 - p x_0 \sum_{j+k=p} x_j x_k \cong 0 \ (\mathrm{mod}\, p).$$

Since $\deg m = q^{2p+6}t^{2p+1}$, the dimension of the kernel of $d_2 : C(2p+6, 2p+1) \to C(2p+6, 2p)$ jumps by 1 when we reduce it modulo p. Therefore its cokernel has $\mathbb{Z}_p$-torsion. $\blacksquare$

We have verified the presence of this torsion in Khovanov homology for $p = 5, 7$.

3 Algebraic structure

We now consider the rational homology of the chain complex appearing in Conjecture 1. We will work with rational coefficients for the remainder of the paper.

3.1 Koszul model

Conjecture 1 tells us to consider the polynomial ring in even variables $x_0, x_1, \ldots, x_{n-1}$ and equal number of odd variables $\xi_0, \xi_1, \ldots, \xi_{n-1}$ bigraded as

$$\deg(x_k) = q^{2k+2}t^{2k}, \quad \deg(\xi_i) = q^{2i+4}t^{2i+1}.$$

The differential d_2 is given by the equation

$$d_2(\xi_m) = \sum_{k=0}^{m} x_k x_{m-k}. \tag{5}$$

One can check that this differential preserves the q-grading and decreases the t-grading by 1.

Remark 12 *Consider the generating functions*

$$x(z) = \sum_{k=0}^{n-1} x_k z^k, \quad \xi(z) = \sum_{k=0}^{n-1} \xi_k z^k.$$

If we work over the ring $\mathbb{Z}[z]/(z^n)$, we can express the differential as

$$d_2(\xi(z)) = x(z)^2.$$

At the bottom level of the Koszul complex, we get the quotient of the polynomial ring in even variables by the ideal I_n generated by the coefficients of the series $x(z)^2$. It was remarked by B. Feigin and A. Stoyanovsky ([13]) that such ideal in the limit $n \to \infty$ corresponds to the integrable representation of $\widehat{sl_2}$ at level 1, and the equation $x(t)^2$ is an example of the Lepowsky-Primc equations ([29]). The bigraded Hilbert series of $\mathbb{C}[x_0, x_1, \ldots]/I_\infty$ was computed in [13] by two different methods, and the identity of the answers corresponds to the following generalization of the Rogers-Ramanujan identity (cf. [2]):

$$H_{q,t}(\mathbb{C}[x_0, x_1, \ldots]/I_\infty) = \sum_{p=0}^{\infty} \frac{q^{2p^2}t^{2p(p-1)}}{(1-q^2t^2)(1-q^4t^4)\ldots(1-q^{2p}t^{2p})} = \tag{6}$$

$$\frac{1}{\prod_{k=1}^{\infty}(1-q^{2k}t^{2k-2})}\sum_{n=0}^{\infty}(-1)^n\prod_{k=1}^{n}\frac{(1-q^{2k}t^{2k-2})}{(1-q^{2k}t^{2k})}(q^{5n^2+n}t^{5n^2-3n}-q^{(n+1)(5n+4)}t^{5n^2+5n})$$

A similar problem was independently studied by C. Brushek, H. Mourtada and J. Schepers in [7], where it appeared in the computation of the Hilbert-Poincaré series of the arc space of double point.

Most of the algebraic constructions below can be considered as a straightforward generalization of these results to the full Koszul homology. In particular, we conjecture the identity (17) that degenerates to (6) at $a = 0$, and holds up to q^{100} in computer experiments.

It is important to note that if we were to replace $d_2(\xi_k)$ with $d_2'(\xi_k) = \sum \alpha_{ik}x_ix_{k-i}$, then for a generic choice of α_{ik} we would have different ranks of homology groups. In example 19 we show that the ranks of the homology cannot be obtained from simple dimension count. As a result, for the random choice of α_{ik} we would have less homology, and, for example, Theorem 35 about the convergence of the Lee's spectral sequence would fail.

3.2 Examples

Example 13 *Let us compute the stable Khovanov homology for* $(2,\infty)$ *case. We have two even generators* x_0, x_1 *and two odd generators* ξ_0, ξ_1*. Since* $d_2(\xi_0) = x_0^2, d_2(\xi_1) = 2x_0x_1$*, we have a non-trivial homology generator* $\mu_0 = 2x_1\xi_0 - x_0\xi_1$*. The stable Khovanov homology is spanned by the elements*

$$p(x_1) + \alpha x_0 + r(x_1)\mu_0$$

(remark that $x_0\mu_0 = d_2(\xi_0\xi_1)$*). The Poincaré series has a form*

$$P_{2,\infty}(q,t) = \frac{(1+q^8t^3)}{(1-q^4t^2)} + q^2.$$

Example 14 *Let us compute the stable unreduced Khovanov homology for* $(3,\infty)$ *case. We add the variables* x_2 *and* ξ_2 *with the differential* $d_2(\xi_2) = 2x_0x_2 + x_1^2$*. Let us compute the homology of* d_2 *in this case.*

Lemma 15 *Let*

$$\mu_0 = 2x_1\xi_0 - x_0\xi_1, \quad \mu_1 = 2x_0\xi_2 - x_1\xi_1 - 4x_2\xi_0.$$

Then

$$d_2(\mu_0) = d_2(\mu_1) = 0, \quad x_0\mu_0 = d_2(\xi_0\xi_1), \quad x_0\mu_1 - x_1\mu_0 = 2d_2(\xi_0\xi_2), \tag{7}$$
$$2x_2\mu_0 + x_1\mu_1 = d_2(\xi_1\xi_2), \quad \mu_1\mu_2 = -2d_2(\xi_0\xi_1\xi_2).$$

One can check that the homology is generated by μ_0 *and* μ_1 *and (7) is the complete set of relations between them (this is a special case of Conjectures 23 and 26 below).*

Lemma 16 *The homology of* d_2 *is spanned by the elements of the form*

$$p_1(x_2) + x_0p_2(x_2) + x_1p_2(x_2) + \alpha\mu_0 + \mu_1(q_1(x_2) + x_0q_2(x_2) + x_1q_2(x_2))$$

Proof. *Modulo the image of d_2 we can eliminate all monomials containing*

$$x_0^2, x_0x_1, x_1^2, x_0\mu_0, x_1\mu_0, x_2\mu_0, \mu_1\mu_2.$$

After this modification the remaining monomials will be linearly independent in the homology. ■

Corollary 17

$$P_{3,\infty}(q,t) = \frac{(1+q^{10}t^5)(1+q^2+q^4t^2)}{1-q^6t^4} + q^8t^3.$$

3.3 Generators and relations

Let us describe the generators in the homology generalizing μ_0 and μ_1 constructed in the previous section.

Lemma 18 *Consider the set of indeterminates $\varepsilon_{a,b}$, where $a+b=r$ is fixed, and $a,b \geq 0$. The system of linear equations*

$$\varepsilon_{a,b+c} + \varepsilon_{b,a+c} + \varepsilon_{c,a+b} = 0 \tag{8}$$

has a nontrivial solution.

Remark that the equations in this system are labelled by triples of integers while the variables are labelled by pairs. Therefore the number of equations is asymptotically quadratic in r, while the number of variables is asymptotically linear, and the system is over-determined.

Proof. Let $\varepsilon_{a,b} = 2a-b$. Then

$$\varepsilon_{a,b+c} + \varepsilon_{b,a+c} + \varepsilon_{c,a+b} = (2a-b-c)+(2b-a-c)+(2c-a-b) = 0.$$

■

Example 19 *Consider the case $r=6$. The system has 7 equations for 7 variables:*

$$2\varepsilon_{0,6} + \varepsilon_{6,0} = 0, \quad \varepsilon_{0,6} + \varepsilon_{1,5} + \varepsilon_{5,1} = 0,$$

$$\varepsilon_{0,6} + \varepsilon_{2,4} + \varepsilon_{4,2} = 0, \quad \varepsilon_{0,6} + 2\varepsilon_{3,3} = 0,$$

$$2\varepsilon_{1,5} + \varepsilon_{4,2} = 0, \quad \varepsilon_{1,5} + \varepsilon_{2,4} + \varepsilon_{3,3} = 0, \quad \varepsilon_{2,4} = 0.$$

Surprisingly, it is has rank 6 and its solution is

$$(\varepsilon_{0,6}, \varepsilon_{1,5}, \varepsilon_{2,4}, \varepsilon_{3,3}, \varepsilon_{4,2}, \varepsilon_{5,1}, \varepsilon_{6,0}) = (-6,-3,0,3,6,9,12).$$

We are ready to present some non-trivial classes in stable homology.

Lemma 20 *Let*

$$\mu_s = \sum_{k=0}^{s+1} \varepsilon_{k,s+1-k} x_k \xi_{s+1-k},$$

where the coefficients $\varepsilon_{a,b}$ are defined by the equation (8). Then $d_2(\mu_s)=0$.

Remark 21 *Since the coefficients of d_2 are quadratic in x-variables, the elements μ_s do not belong to the image of d_2.*

Proof. We have to check that

$$d_2(\mu_s) = \sum_{k=1}^{s+2} \varepsilon_{k,s+3-k} x_k d_2(\xi_{s+3-k}) = 0.$$

Let us expand $d_2(\xi_{s+3-k})$:

$$d_2(\mu_s) = \sum_{k=1}^{s+2} \varepsilon_{k,s+3-k} x_k \sum_{j=1}^{k-1} x_j x_{s+3-k-j} = \sum_{i+j+k=s+3} (\varepsilon_{i,j+k} + \varepsilon_{j,i+k} + \varepsilon_{k,i+j}).$$

It rests to apply the equation (8). ■

Remark 22 *Example 19 shows that for a generic choice of the coefficients of d_2 the corresponding 7×7 matrix would be non-degenerate, and the d_2 homology would have* smaller *dimension. In particular, for a generic choice of the coefficients d_2 would have no homology in bidegree $q^{18}t^{13}$. Indeed, the only monomials in this bidegree are $x_a \xi_b$ with $a + b = 6$. The differential maps the space they span to the space spanned by monomials of the form $x_a x_b x_c$ with $a + b + c = 6$ according to the matrix from Example 19.*

Conjecture 23 *The homology of d_2 is generated as an algebra by μ_s and x_i.*

Remark 24 *In what follows we will use the following description of the generators μ_s. Consider the generating function $\mu(z) = \sum_{s=0}^{\infty} z^s \mu_s$. Then*

$$\mu(z) = 2\dot{x}(z)\xi(z) - x(z)\dot{\xi}(z). \tag{9}$$

Let us describe the generalization of the relations (15).

Lemma 25 *The following relations hold in the homology of d_2:*

$$x(z)^2 = 0, \quad x(z)\mu(z) = 0, \quad \ddot{x}(z)\mu(z) - \dot{x}(z)\dot{\mu}(z) = 0, \quad \mu(z)\dot{\mu}(z) = 0. \tag{10}$$

As before, these relations are to be interpreted as holding modulo z^n.

Proof.

$$x\mu = 2x\dot{x}\xi - x^2\dot{\xi} = d_2(\xi\dot{\xi})$$

$$\ddot{x}\mu - \dot{x}\dot{\mu} = \ddot{x}(2\dot{x}\xi - x\dot{\xi}) - \dot{x}(2\ddot{x}\xi + 2\dot{x}\dot{\xi} - \dot{x}\dot{\xi} - x\ddot{\xi}) =$$

$$-x\ddot{x}\dot{\xi} - \dot{x}^2\dot{\xi} + x\dot{x}\ddot{\xi} = -\frac{1}{2}d_2(\dot{\xi}\ddot{\xi}).$$

$$\mu\dot{\mu} = (2\dot{x}\xi - x\dot{\xi})(2\ddot{x}\xi + \dot{x}\dot{\xi} - x\ddot{\xi}) =$$

$$2\dot{x}^2\xi\dot{\xi} - 2x\dot{x}\xi\ddot{\xi} + 2x\ddot{x}\xi\dot{\xi} + x^2\dot{\xi}\ddot{\xi} = d_2(\xi\dot{\xi}\ddot{\xi}).$$

■

Conjecture 26 *The ideal of relations in the Koszul homology is generated by the coefficients of the relations (10).*

Using `Singular` [8], we have verified that both conjectures hold for $n \leq 7$. Some further evidence for the conjectures is provided by Theorem 46 in the next section.

Remark 27 *Remark that the relations (10) are not independent, and there are lots of syzygies between them. For example, first equation presents $d_2(\xi\dot{\xi})$ in terms of $\mu' s$, so its derivative presents $d_2(\xi\ddot{\xi})$ in terms of μ's. On the other hand, the second equation presents $d_2(\dot{\xi}\ddot{\xi})$ in terms of μ's. This suggests a syzygy*

$$d_2(\xi\dot{\xi})d_2(\ddot{\xi}) - d_2(\xi\ddot{\xi})d_2(\dot{\xi}) + d_2(\dot{\xi}\ddot{\xi})d_2(\xi) = d_2^2(\xi\dot{\xi}\ddot{\xi}) = 0.$$

Lemma 28 *Assuming Conjecture 23, $\mathrm{Kh}_{alg}(n,\infty)$ contains at most $\lfloor\frac{n+1}{3}\rfloor$ "levels", i. e. the maximal ξ–degree of a homology generator is at most $\lfloor\frac{n+1}{3}\rfloor$.*

Proof. Consider the equation $\mu(z)\dot{\mu}(z) = 0$. For the odd powers of z, the coefficients look like $\mu_i\mu_{i+1} + \ldots = 0$, for the even powers of z, the coefficients look like $\mu_i\mu_{i+2} + \ldots = 0$ (recall that μ_i are odd, so there are no terms $\mu_i\mu_i$).

Therefore one can eliminate all monomials containing $\mu_i\mu_{i+1}$ and $\mu_i\mu_{i+2}$, and the monomial of the maximal ξ-degree is $\mu_0\mu_3\cdots\mu_{\lfloor\frac{n-2}{3}\rfloor}$. Its degree is $\lfloor\frac{n-2}{3}\rfloor + 1 = \lfloor\frac{n+1}{3}\rfloor$. ■

Finally, let us explain some corollaries of the recent work by Feigin [12]. His work provides further evidences for the above mentioned conjectures. Feigin studies properties of the ideal J inside $R = \mathbb{C}[\xi_0, \xi_1, \ldots, x_0, x_1, \ldots]$ generated by coefficients of the power series:

$$\mu(z), \quad \xi(z)\dot{\xi}(z), \quad \xi(z)\ddot{\xi}(z), \quad \xi(z)\dddot{\xi}(z).$$

An easy computation shows that J is preserved by d_2; the main object of study of [12] is the differential graded algebra $R_{(1)} = R/J$. Feigin shows that $R_{(1)}$ is a representation of the Virasoro algebra generated by L_i, $i \in \mathbb{Z}$ and c:

$$[L_i, c] = 0, \quad [L_m, L_n] = (m-n)L_{m+n} + \delta_{m+n}\frac{m^3 - m}{3}c.$$

The central element c acts by the constant $-4/5$ on $R_{(1)}$. The algebra $R_{(1)}$ has a natural grading by the odd variables:

$$\deg(\xi_i) = 1, \quad \deg(x_i) = 0.$$

The graded components $R_{(1)}[j]$ are subrepresentations and Feigin identifies them with some particular highest weight modules of L. Let us briefly recall the basics of the highest weight theory for the Virasoro algebra.

The algebra L naturally splits into three parts: the positive part L^+ generated by L_i, $i > 0$; the negative part L^- generated by L_i, $i < 0$ and the span of L_0, c. From the relations for L we see that L_0 is a grading operator; it is customary to call the eigenspaces of L_0 levels. The vector from a L-module is called singular if it is anihilated by L^+. The Verma module M_λ is a L-module freely generated from the singular vector on the level λ.

Theorem 29 *[12] We have*

1. *The differential* d_2 *commutes with the action of the Virasoro algebra.*
2. *The graded component* $R_{(1)}[j]$ *is isomorphic to the quotient of the Verma module* M_{λ_i}, $\lambda_i = (5i^2 - 3i)/2$ *by a singular vector at the level* $2i + 1$.
3. *For any* $i > 0$, $H^i(R_{(1)}, d_2) = 0$
4. $H^0(R_{(1)}, d_2)$ *is the irreducible quotient of* M_0.

Corollary 30 *Lower level of* $\mathrm{Kh}_{alg}(\infty, \infty)$ *is the irreducible representation of the Virasoro algebra.*

All generators besides of the ideal J besides μ_i have odd degree at least 2 hence

Corollary 31 *The first homology of* d_2 *is generated by* μ_i *and* x_i.

Remark 32 *The theorem above is just a particular case of more general result from [12] that might be relevant for studies of* $sl(N)$ *homology. Also it is very plausable that one can extend the result of the last corollary to the higher homological degrees by some bootstrapping procedure. We hope to return to this question in our future work.*

3.4 Lee's spectral sequence

It was conjectured in [9] that Lee's spectral sequence ([28],[34]) is induced by a differential d_1 that commutes with d_2. We propose a formula for this differential:

$$d_1(\xi_i) = x_i.$$

Remark that if d_1 satisfies the Leibnitz rule, it is uniquely defined by the grading restrictions.

Let us consider the spectral sequence induced by d_1:

Example 33 *Consider the unreduced stable homology of the* $(2, \infty)$ *torus knot. As it was shown in the Example 13, the homology is generated by* x_0, x_1 *and* μ_0 *modulo relations*

$$x_0^2 = 2x_0x_1 = x_0\mu_0 = 0.$$

Remark that

$$d_1(\mu_0) = d_1(2x_1\xi_0 - x_0\xi_1) = 2x_1x_0 - x_0x_1 = x_0x_1 = \frac{1}{2}d_2(\xi_1).$$

This means that the second differential in the spectral sequence (knight move) acts as

$$\delta(\mu_0) = d_1 \circ d_2^{-1} \circ d_1(\mu_0) = d_1(\frac{1}{2}\xi_1) = \frac{1}{2}x_1.$$

Therefore μ_0 *kills* ξ_1 *by the knight move, and the spectral sequence converges at* E_3 *page to the two-dimensional space*

$$E_3 = E_\infty =< 1, x_0 > .$$

Remark 34 *One can find an apparent contradiction in this result — the homology of d_1 is clearly one-dimensional, while the spectral sequence converges to a two-dimensional space. This problem is caused by the infinite dimension of the homology. One can check that for a finite $(2,m)$ knot (i.e. for a suitable finite-dimensional quotient of this complex) the homology of d_1 will be two-dimensional — one generator will be 1, while the degree of the second one will grow as m increases.*

Motivated by this example, we formulate the following algebraic counterpart for the conjectures from [5] and [34]. The following theorem holds module Conjectures 23 and 26.

Theorem 35 *Consider the spectral sequence induced by d_1 acting in the homology of d_2. Then*
a) $E_1 = E_2 = H^*(d_2)$.
b) $E_3 = E_\infty =< 1, x_0 >$.
In particular, the spectral sequence converges at the E_3 page.

Proof. We know that the stable homology are generated by μ_s and x_i, therefore the multiplicativity of the spectral sequence allows us to focus on these generators. Remark that

$$d_1(\mu_s) = \sum_{k=0}^{s+1} \varepsilon_{k,s+1-k} x_k x_{s+1-k} = \frac{s+1}{2} \sum_{k=0}^{s+1} x_k x_{s+1-k} = \frac{s+1}{2} d_2(\xi_{s+1}). \qquad (11)$$

Here we used the equation

$$\varepsilon_{k,s+1-k} + \varepsilon_{s+1-k,k} = 2k - (s+1-k) + 2(s+1-k) - k = s+1.$$

We can compute the second differential in the spectral sequence $\delta(\mu_s)$ using the equation (11):

$$\delta(\mu_s) = d_1 \circ d_2^{-1} \circ d_1(\mu_s) = d_1(\frac{s+1}{2}\xi_{s+1}) = \frac{s+1}{2} x_{s+1}.$$

Therefore all even generators x_i will be killed by μ_{i-1} by the knight move. ∎

Remark 36 *One can reformulate this proof in terms of the generating series. We have*

$$d_1(\mu(z)) = d_1(-x(z)\dot{\xi}(z) + 2\dot{x}(z)\xi(z)) = -x(z)\dot{x}(z) + 2x(z)\dot{x}(z) = x(z)\dot{x}(z) = \frac{1}{2} d_2(\dot{\xi}(z)).$$

Therefore

$$\delta(\mu(z)) = d_1 \circ d_2^{-1} \circ d_1(\mu(z)) = d_1(\frac{1}{2}\dot{\xi}(z)) = \frac{1}{2}\dot{x}(z).$$

One can prove the similar theorem for the reduced homology (see below). In the reduced case, the E_∞ term will be one-dimensional and spanned by 1.

4 Poincaré polynomials

4.1 Bosonic formula

In this section we give a conjectural formula for the Poincare polynomial of $\mathrm{Kh}_{alg}(n,\infty)$ for all n. This formula comes from the computer experiments, and it can be considered as a potential generalization of the "bosonic" side of the Rogers-Ramanujan identity in [13] (see also [22], [14], [30]). It is worth to note that this "bosonic formula" was obtained in [13] with the usage of the localization technique on the affine flag variety for $\widehat{sl(2)}$. We plan to compare this approach with the one proposed below in the future.

Recall that the lower level of the stable unreduced Khovanov homology of (n,∞) knot can be described by the quotient of $\mathbb{C}[x_0,\ldots,x_{n-1}]$ by the ideal generated by the coefficients of $x(z)^2$ at $z^0,\ldots,z^{n-1}$.

Conjecture 37 *Let $z=q^2t^2$. The unreduced Hilbert series for the lower level of $\mathrm{Kh}_{alg}(n,\infty)$ has the form*

$$L_n(q,t)=\frac{1}{\prod_{k=1}^{n}(1-q^{2k}t^{2k-2})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}(1-q^{2k}t^{2k-2})\times \tag{12}$$

$$\left(q^{5p^2+p}t^{5p^2-3p}\binom{n-2p+1}{p}_z-q^{(p+1)(5p+4)}t^{5p^2+5p}\binom{n-2p-1}{p}_z\right).$$

Here we use the standard z-binomial notation:

$$[m!]_z=\prod_{k=1}^{m}\frac{1-z^k}{1-z},\quad \binom{m}{l}_z=\frac{[m!]_z}{[l!]_z[m-l!]_z}\quad (m\geq l).$$

Remark 38 *In the limit $n\to\infty$ the z-binomial coefficients degenerate to simple products:*

$$\binom{n-2p+1}{p}_z,\binom{n-2p-1}{p}_z\xrightarrow{n\to\infty}\frac{1}{(1-z)(1-z^2)\ldots(1-z^p)}=\prod_{k=1}^{p}\frac{1}{(1-q^{2k}t^{2k})},$$

therefore the equation (12) has a limit

$$L_\infty(q,t)=\frac{1}{\prod_{k=1}^{n}(1-q^{2k}t^{2k-2})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}\frac{(1-q^{2k}t^{2k-2})}{(1-q^{2k}t^{2k})}\left(q^{5p^2+p}t^{5p^2-3p}-q^{(p+1)(5p+4)}t^{5p^2+5p}\right).$$

This is the right hand side of the generalized Rogers-Ramanijan identity (6), and therefore in this limit the Conjecture 37 follows from the results of [13].

One can try to extend the equation (12) to higher levels of the Koszul homology.

Conjecture 39 *The unreduced Hilbert series for* $\mathrm{Kh}_{alg}(n,\infty)$ *has the form*

$$P_n(q,t) = \frac{1}{\prod_{k=1}^{n}(1-q^{2k}t^{2k-2})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}(1-q^{2k}t^{2k-2}) \times \tag{13}$$

$$\times \prod_{k=3p+1}^{n-1}(1+q^{2k+6}t^{2k+1})\prod_{k=1}^{2p-1}(1+q^{2k+2}t^{2k-1})\times$$

$$[q^{5p^2+p}t^{5p^2-3p}(1+\chi_p^+ q^{6p+4}t^{6p-1})(1+q^{6p+6}t^{6p+1})\binom{n-2p+1}{p}_z +$$

$$q^{5p^2+7p+2}t^{5p^2+3p-1}(1+q^{6p+6}t^{6p+1})(1-q^{2p+2}t^{2p})\binom{n-2p}{p}_z -$$

$$q^{5p^2+9p+4}t^{5p^2+5p}(1+q^{2p+2}t^{2p+1})(1+\chi_p^+ q^{4p+2}t^{4p-1})\binom{n-2p-1}{p}_z],$$

where $\chi_p^+ = 0$ *when* $p=0$, $\chi_p^+ = 1$ *for* $p>0$ *and the second product under inside sum is* 1 *when* $3p+1 > n-1$.

4.2 Fermionic formula for $T(\infty,\infty)$

Based on the experimental data, A. Shumakovich and P. Turner conjectured that the Poincaré polynomials for the Khovanov homology for $(n,n+1)$ torus knots satisfy the following recurrence relation.

Conjecture 40 *([37]) Let* $\mathcal{K}_n(q,t)$ *denote the Poincaré polynomials for the Khovanov homology for the* $(n,n+1)$ *torus knot. Then*

$$\mathcal{K}_n(q,t) = \mathcal{K}_{n-1}(q,t) + \mathcal{K}_{n-2}(q,t)q^{2n}t^{2n-2} + \mathcal{K}_{n-3}(q,t)q^{2n+4}t^{2n-1}. \tag{14}$$

Let us construct a combinatorial model for this recursion relation. Consider the sequences of 0's and 1's with no blocks of the form 1111 anywhere and no blocks 111 except, possibly, the beginning. Such sequences are split (outside the beginning) into 1's and 11's separated by blocks of 0's.

Example 41 *For* $n=3$ *all 8 sequences are admissible. For* $n=4$ *we have 14 sequences:* 1111 *and* 0111 *are forbidden.*

We weight such sequences by a product of terms corresponding to blocks of 1's appearing in the sequence. The weights are as follows:

1) 111 in the beginning: $q^{12}t^5$;

2) 1 at position n (first digit has position 0): $q^{2n+2}t^{2n}$;

3) 11 starting at position n: $q^{2n+8}t^{2n+3}$.

Let K_n be the weighted state sum for length n sequences; that is, the sum of the weights for all such sequences.

Lemma 42 *K_n satisfies the recursion relation* (14) *and agrees with $\mathcal{K}_n$ for $n = 1, 2, 3$.*

Proof. Let us check the recursion relation. If a sequence ends with 0, this accounts to K_{n-1}. If a sequence ends with 01, this accounts to $K_{n-2}(q,t)q^{2n}t^{2n-2}$. If a sequence ends with 011, this accounts to $K_{n-3}(q,t)q^{2n+4}t^{2n-1}$. The values of K_n for $n = 1, 2, 3$ are easily checked. ■

Let us write the formula for the limit $K(q,t) = \lim_{n\to\infty} K_n(q,t)$.

Theorem 43

$$K(q,t) = \sum_{p=0}^{\infty} q^{2p^2} t^{2p(p-1)} (1 + q^{8p+12}t^{8p+5}) \frac{(1+q^6t^3)(1+q^8t^5)\dots(1+q^{2p+4}t^{2p+1})}{(1-q^2t^2)(1-q^4t^4)\dots(1-q^{2p}t^{2p})}. \quad (15)$$

Proof. First, let us assume that we have no 111 in the beginning. Let $U_p(q,t)$ be the state sum with p blocks of units. A sequence with $p+1$ blocks can be one of the following:

1) Starting with 10 at position k, what accounts to $q^{2k+2}t^{2k} \cdot (q^2t^2)^{p(k+2)} U_p$. If we sum over all k, we get

$$\frac{q^{4p+2}t^{4p}}{1 - q^{2(p+1)}t^{2(p+1)}} U_p.$$

2) Starting with 110 at position k, what accounts to $q^{2k+8}t^{2k+3} \cdot (q^2t^2)^{p(k+3)} U_p$ If we sum over all k, we get

$$\frac{q^{6p+8}t^{6p+3}}{1 - q^{2(p+1)}t^{2(p+1)}} U_p.$$

Summing up, we get

$$\frac{q^{4p+2}t^{4p}(1 + q^{2p+6}t^{2p+3})}{1 - q^{2(p+1)}t^{2(p+1)}} U_p.$$

Therefore

$$U_p(q,t) = q^{2p^2}t^{2p(p-1)} \frac{(1+q^6t^3)(1+q^8t^5)\dots(1+q^{2p+4}t^{2p+1})}{(1-q^2t^2)(1-q^4t^4)\dots(1-q^{2p}t^{2p})}.$$

Let $V_p(q,t)$ denote the state sum where we allow 111 in the beginning. Then

$$V_p = U_p + q^{12}t^5(q^2t^2)^{4p}U_p = (1 + q^{8p+12}t^{8p+5})U_p.$$

■

We can verify directly that the Euler characteristic of $K(q,t)$ agrees with the stable Jones polynomial of $T(\infty,\infty)$:

Lemma 44

$$K(q,-1) = \frac{1}{1-q^2}.$$

Proof. Remark that

$$U_p(q,-1)=q^{2p^2}\frac{(1-q^6)(1-q^8)\dots(1-q^{2p+4})}{(1-q^2)(1-q^4)\dots(1-q^{2p})}=q^{2p^2}\frac{(1-q^{2p+2})(1-q^{2p+4})}{(1-q^2)(1-q^4)},$$

$$V_p(q,-1)=(1-q^{8p+12})U_p(q,-1).$$

Therefore we have to prove that

$$\sum_{p=0}^{\infty} q^{2p^2}(1-q^{2p+2})(1-q^{2p+4})(1-q^{8p+12})=1-q^4. \tag{16}$$

This follows from the direct expansion of the left hand side: all terms will cancel out except 1 and $-q^4$. ■

Comparing the "fermionic" formula (15)) with the "bosonic" formula (13) in the limit $n\to\infty$ suggests the following identity. The a-grading from HOMFLY homology can be traced on both sides.

Conjecture 45 *("Khovanov-Rogers-Ramanujan identity") Let*

$$A(a,q,t)=\sum_{p=0}^{\infty} q^{2p^2}t^{2p(p-1)}(1+a^2q^{8p+8}t^{8p+5})\prod_{j=1}^{p}\frac{(1+a^2q^{2j}t^{2j+1})}{(1-q^{2j}t^{2j})}.$$

and

$$B(a,q,t)=\frac{1}{\prod_{k=1}^{\infty}(1-q^{2k}t^{2k-2})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}\frac{(1-q^{2k}t^{2k-2})}{(1-q^{2k}t^{2k})}\times$$
$$\times\prod_{k=3p+1}^{\infty}(1+a^2q^{2k+2}t^{2k+1})\prod_{k=1}^{2p-1}(1+a^2q^{2k-2}t^{2k-1})\times$$
$$[q^{5p^2+p}t^{5p^2-3p}(1+\chi_p^{+}a^2q^{6p}t^{6p-1})(1+a^2q^{6p+2}t^{6p+1})+$$
$$a^2q^{5p^2+7p-2}t^{5p^2+3p-1}(1+a^2q^{6p+2}t^{6p+1})(1-q^{2p+2}t^{2p})-$$
$$q^{5p^2+9p+4}t^{5p^2+5p}(1+a^2q^{2p-2}t^{2p+1})(1+\chi_p^{+}a^2q^{4p-2}t^{4p-1})].$$

Then

$$A(a,q,t)=B(a,q,t). \tag{17}$$

Using a computer, we have checked that the identity holds up through terms of order q^{100}.

The following theorem provides some evidence in support of Conjectures 23 and 26.

Theorem 46 *The Hilbert series of the algebra generated by μ_n ($n\geq 0$) and x_i ($i\geq 0$) and satisfying the relations in equation* (10) *is $K(q,t)$.*

Proof. The elements μ_n have grading

$$\deg(\mu_n) = q^{2n+8}t^{2n+3}.$$

Let us return to our combinatorial model:

1) 1 at position n corresponds to x_n.
2) 11 starting at position n corresponds to μ_n.
3) 111 in the beginning corresponds to $x_0\mu_1$.

We have to check that we can eliminate the following products using the relations:

$$x_i x_{i+1}, \quad x_i\mu_{i+1} \quad (i > 0), \quad \mu_i x_{i+2}, \quad \mu_i\mu_{i+2},$$
$$x_i^2, \quad x_i\mu_i, \quad x_{i+1}\mu_i, \quad \mu_i\mu_{i+1}.$$

We can eliminate x_i^2 and $x_i x_{i+1}$ using the equation $x(z)^2 = 0$; $\mu_i\mu_{i+1}$ and $\mu_i\mu_{i+2}$ using the equation $\mu(t)\dot{\mu}(z) = 0$. Finally, we can eliminate $x_{i-1}\mu_{i+1}, x_i\mu_i, x_{i+1}\mu_i, x_{i+2}\mu_i$ using two remaining equations

$$x(z)\mu(z) = \ddot{x}(z)\mu(z) - \dot{x}(z)\dot{\mu}(z) = 0.$$

■

5 Reduced homology

In this section we briefly review the structure of the reduced stable homology. We conjecture that the construction and the differential are exactly the same as in the unreduced case, but we have to omit x_0 and ξ_0. Therefore we get the following result.

Consider the polynomial ring in even variables $x_1, \ldots, x_{n-1}$ and equal number of odd variables $\xi_1, \ldots, \xi_{n-1}$ bigraded as

$$\deg(x_k) = q^{2k+2}t^{2k}, \quad \deg(\xi_i) = q^{2i+4}t^{2i+1}.$$

The differential d_2 is given by the equation

$$d_2(\xi_m) = \sum_{k=1}^{m-1} x_k x_{m-k}. \tag{18}$$

Conjecture 47 *The stable reduced Khovanov homology of (n, ∞) knot is isomorphic to the homology of $\mathbb{C}[x_1 \ldots, x_{n-1}, \xi_1, \ldots, \xi_{n-1}]$ with respect to d_2.*

Example 48 *(cf. [16]) Let us compute the reduced homology of the $(4, \infty)$ knot. We have*

$$d_2(\xi_1) = 0, \quad d_2(\xi_2) = x_1^2, \quad d_2(\xi_3) = 2x_1x_2.$$

Analogously to the Example 13, we can introduce a homology generator $\overline{\mu_0} = 2x_2\xi_2 - x_1\xi_3$ and check that the homology is spanned by the expressions of the form $p(x_2) + \alpha x_1 + \overline{\mu_0}q(x_2)$ up to multiplication by polynomials in ξ_1 and x_3. Therefore the Poincaré series has a form

$$\overline{P}_{4,\infty}(q,t) = \frac{(1+q^6t^3)}{(1-q^8t^6)}\left(q^4t^2 + \frac{1+q^8t^5}{1-q^6t^4}\right).$$

Remark that the reduced and unreduced differentials look similar up to a shift of grading. More precisely, modulo the multiplication by ξ_1 and x_{n-1} we can replace ξ_i by ξ_{i-2} and x_i by x_{i-1} to get the unreduced stable homology of the $(n-2,\infty)$ knot. We get the following result.

Lemma 49 *Modulo the multiplication by ξ_1 and x_{n-1} the reduced stable homology of the (n,∞) torus knot is isomorphic (as a vector space) to the unreduced stable homology of the $(n-2,\infty)$ torus knot.*

One can reconstruct the grading shifts for this correspondence, and obtain an analogue of the equation (13) for the reduced homology.

Conjecture 50 *The reduced Hilbert series for the (n,∞) knot Khovanov homology has a form*

$$\overline{P}_n(q,t)=\frac{(1+q^6t^3)}{\prod_{k=1}^{n-1}(1-q^{2k+2}t^{2k})}\sum_{p=0}^{\infty}(-1)^p\prod_{k=1}^{p}(1-q^{2k+2}t^{2k})\times \tag{19}$$

$$\times\prod_{k=3p+1}^{n-1}(1+q^{2k+12}t^{2k+7})\prod_{k=1}^{2p-1}(1+q^{2k+6}t^{2k+3})\times$$

$$[q^{5p^2+5p}t^{5p^2+p}(1+q^{6p+10}t^{6p+5})(1+q^{6p+12}t^{6p+7})\binom{n-2p+1}{p}_z+$$

$$q^{5p^2+11p+6}t^{5p^2+7p+3}(1+q^{6p+12}t^{6p+7})(1-q^{2p+4}t^{2p+2})\binom{n-2p}{p}_z-$$

$$q^{5p^2+13p+8}t^{5p^2+9p+4}(1+q^{2p+4}t^{2p+3})(1+q^{4p+6}t^{4p+3})\binom{n-2p-1}{p}_z],$$

where the second product under inside sum is 1 when $3p+1>n-1$.

Remark 51 *It was conjectured in [9] that*

$$\overline{P}_{2,\infty}(q,t)=(1+q^6t^3)(1+q^4t^2+q^8t^4+\ldots),$$

$$\overline{P}_{3,\infty}(q,t)=\frac{1+q^4t^2+q^6t^3+q^{10}t^5}{1-q^6t^4},$$

$$\overline{P}_{4,\infty}(q,t)=\frac{(1+q^6t^3)}{(1-q^8t^6)}\left(1+q^4t^2+\frac{q^6t^4(1+q^8t^5)}{1-q^6t^4}\right).$$

One can check that these answers coincide with the above construction (see Appendix B for a comparison).

6 Matrix factorizations

In different physical models of knot homology (e.g. [18],[19]) the colored homology of the unknot is described in terms of the matrix factorizations. Following the discussions in previous section, we are focusing on the unknot coloured by the nth symmetric power of the fundamental representation of $\mathfrak{sl}(2)$.

Definition 52 *([11]) A matrix factorisation of a function W over a ring R is a pair (M, d), where $M = M_0 + M_1$ is a $\mathbb{Z}_2$-graded R-modules of finite rank equipped with an R-linear map d of odd degree satisfying the equation $d^2 = W \cdot id_M$.*

We will need the following basic facts about matrix factorizations:

Theorem 53 *(a) ([20]) The Hochschild cohomology of the algebra of functions on $\mathbb{C}^n$ equals to the algebra of polyvector fields on $\mathbb{C}^n$.*

(b) ([10]) Consider a function $W : \mathbb{C}^n \to \mathbb{C}$. Then the Hochschild cohomology $HH(MF(W))$ of the category of matrix factorizations of W equals the Koszul homology of the complex obtained from polyvector fields by the contraction with dW.

(c) (e.g. [10]) If W has an isolated singularity, then $HH(MF(W))$ is isomorphic to the Milnor algebra of W at this singularity:

$$HH(MF(W)) = \mathbb{C}[x_1, \ldots, x_n] / \left(\frac{\partial W}{\partial x_1}, \ldots, \frac{\partial W}{\partial x_n} \right).$$

Remark that we can dualize the Koszul complex (b) and obtain the differential

$$D_W(dx_i) = \frac{\partial W}{\partial x_i}$$

acting as a derivation on the algebra $\mathbb{C}[x_1, \ldots, x_n, dx_1, \ldots, dx_n]$ of differential forms on $\mathbb{C}^n$. Now the part (c) follows from the well known fact that W has an isolated singularity if and only if its partial derivatives form a regular sequence.

The following potentials (with isolated singularities) were proposed for the totally symmetric representations by S. Gukov and J. Walcher:

Proposition 54 *([18]) The generating function for the $(\mathfrak{sl}(N), S^k)$ potentials has a form:*

$$\sum_{N=0}^{\infty} z^{N+k} (-1)^N W_{\mathfrak{sl}(N), S^k}(x_1, \ldots, x_k) = \left(1 + \sum_{i=1}^{k} z^i x_i \right) \ln \left(1 + \sum_{i=1}^{k} z^i x_i \right)$$

The$(\mathfrak{sl}(N), S^k)$ colored homology of the unknot coincides with the Milnor algebra of the corresponding potential.

We slightly change notations and write

$$W_{phys}(x_0, \ldots, x_{n-1}) = W_{\mathfrak{sl}(2), S^n}(x_0, \ldots, x_{n-1}) = \text{Coef}_{n+2} \left[(1 + zx(z)) \ln(1 + zx(z)) \right],$$

where $x(z) = \sum_{i=0}^{n-1} z^i x_i$.

Let us assume that x_i are bigraded as above: $\deg(x_i) = q^{2i+2}t^{2i}$. Then the differential D_W will preserve both gradings iff W is bihomogeneous. Let $\overline{W}_n(x_0, \ldots, x_{n-1})$ be the piece of bidegree $(2n+4, 2n-2)$ in W_{phys}.

Lemma 55 *The potential $\overline{W}_n$ can be written as follows:*

$$\overline{W_n}(x_0, \ldots, x_{n-1}) = -\frac{1}{6}\text{Coef}_{n-1}[x(z)^3].$$

Proof. Remark that the difference between q- and t-gradings for x_i equals to 2. Therefore the piece of bidegree $(2n+4, 2n-2)$ should be cubic in x_i. Now

$$(1 + zx(z))\ln(1 + zx(z)) = zx(z) + \frac{1}{2}z^2x(z)^2 - \frac{1}{6}z^3x(z)^3 + \ldots,$$

so the cubic part equals to $-\frac{1}{6}z^3x(z)^3$. ■

Example 56 *We have*

$$\overline{W}_0 = -\frac{1}{6}x_0^3, \quad \overline{W}_1 = -\frac{1}{2}x_0^2x_1.$$

Remark that $\overline{W}_1$ has a non-isolated singularity.

Theorem 57 *The Hochschild homology of the category of matrix factorizations of the potential $\overline{W}_n$ is isomorphic to the homology of d_2.*

Proof. From Theorem 53 it is sufficient to study the Koszul complex associated with the partial derivatives of $\overline{W}_n$. We have

$$\frac{\partial}{\partial x_i}\overline{W}_n = -\frac{1}{6}\frac{\partial}{\partial x_i}\text{Coef}_{n-1}[x(z)^3] = -\frac{1}{2}\text{Coef}_{n-1}[x(z)^2\frac{\partial}{\partial x_i}x(z)] =$$

$$-\frac{1}{2}\text{Coef}_{n-1}[z^ix(z)^2] = -\frac{1}{2}\text{Coef}_{n-1-i}[x(z)^2].$$

Therefore

$$D_{\overline{W}_n}(dx_{n-1-i}) = \frac{\partial}{\partial x_{n-1-i}}\overline{W}_n = -\frac{1}{2}\text{Coef}_i[x(z)^2] = -\frac{1}{2}\sum_{j=0}^{i} x_jx_{j-i}.$$

It rests to identify

$$\xi_i = -2dx_{n-1-i}.$$

■

Appendix

A Unreduced Poincaré series

Here we collect the answers for the Poincaré series of $\mathrm{Kh}_{alg}(n, \infty)$ for $n \leq 7$. These series were computed using `Singular` [8]. The resulting series coincide with Shumakovitch's computations [37] up to high q-degree. For example, for $(7, 20)$ torus knot the first difference is in q-degree 42.

For the reader's convenience, we multiply both parts of the equation (13) by $\prod_{i=1}^{n}(1 - q^{2i}t^{2i-2})$.

$$(1-q^2)(1-q^4t^2)P_2(q,t) = (1+q^8t^3) - q^4(1+q^8t^3) - q^6t^2(1-q^2)(1+q^4t) =$$
$$(1-q^2)(1+q^2-q^6t^2+q^8t^3);$$

$$\prod_{i=1}^{3}(1-q^{2i}t^{2i-2})P_3(q,t) = (1+q^8t^3)(1+q^{10}t^5) - q^4(1+q^8t^3)(1+q^{10}t^5) -$$
$$q^6t^2(1-q^2)(1+q^4t)(1+q^2t^2)(1+q^{10}t^5) - q^{14}t^7(1-q^2)(1-q^4t^2)(1+q^4t) =$$
$$(1-q^2)(1-q^4t^2)(1+q^2+q^4t^2+q^8t^3+q^{10}t^5+q^{12}t^5);$$

$$\prod_{i=1}^{4}(1-q^{2i}t^{2i-2})P_4(q,t) = (1+q^8t^3)(1+q^{10}t^5)(1+q^{12}t^7) - q^4(1+q^8t^3)(1+q^{10}t^5)(1+q^{12}t^7) -$$
$$q^6t^2(1-q^2)(1+q^4t)(1+q^{10}t^5)(1+q^{12}t^7)(1+q^2t^2+q^4t^4) -$$
$$q^{14}t^7(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^{12}t^7)(1+q^2t^2) + q^{18}t^{10}(1-q^2)(1+q^4t)(1+q^6t^3)(1+q^4t^3);$$

$$\prod_{i=1}^{5}(1-q^{2i}t^{2i-2})P_5(q,t) = \prod_{i=1}^{4}(1+q^{2i+6}t^{2i+1}) - q^4\prod_{i=1}^{4}(1+q^{2i+6}t^{2i+1}) -$$
$$q^6t^2(1-q^2)(1+q^4t)(1+q^2t^2+q^4t^4+q^6t^6)\prod_{i=2}^{4}(1+q^{2i+6}t^{2i+1}) -$$
$$q^{14}t^7(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^{12}t^7)(1+q^{14}t^9)(1+q^2t^2+q^4t^4) +$$
$$q^{18}t^{10}(1-q^2)(1+q^4t)(1+q^6t^3)(1+q^2t^2)(1+q^{14}t^9)(1+q^4t^3) +$$
$$q^{22}t^{14}(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^6t^3)(1+q^8t^5);$$

$$\prod_{i=1}^{6}(1-q^{2i}t^{2i-2})P_6(q,t) = \prod_{i=1}^{5}(1+q^{2i+6}t^{2i+1}) - q^4\prod_{i=1}^{5}(1+q^{2i+6}t^{2i+1}) -$$

$$q^6t^2(1-q^2)(1+q^4t)(1+q^2t^2+q^4t^4+q^6t^6+q^8t^8)\prod_{i=2}^{5}(1+q^{2i+6}t^{2i+1})-$$
$$q^{14}t^7(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^{12}t^7)(1+q^{14}t^9)(1+q^{16}t^{11})(1+q^2t^2+q^4t^4+q^6t^6)+$$
$$q^{18}t^{10}(1-q^2)(1+q^4t)(1+q^6t^3)(1+q^2t^2+q^4t^4)(1+q^{14}t^9)(1+q^{16}t^{11})(1+q^4t^3)+$$
$$q^{22}t^{14}(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^6t^3)(1+q^8t^5)(1+q^{16}t^{11})(1+q^2t^2+q^4t^4)+$$
$$q^{36}t^{25}(1-q^2)(1-q^4t^2)(1-q^6t^4)(1+q^4t)(1+q^6t^3)(1+q^8t^5);$$

$$\prod_{i=1}^{7}(1-q^{2i}t^{2i-2})P_7(q,t)=\prod_{i=1}^{6}(1+q^{2i+6}t^{2i+1})-q^4\prod_{i=1}^{6}(1+q^{2i+6}t^{2i+1})-$$
$$q^6t^2(1-q^2)(1+q^4t)(1+q^2t^2+q^4t^4+q^6t^6+q^8t^8+q^{10}t^{10})\prod_{i=2}^{6}(1+q^{2i+6}t^{2i+1})-$$
$$q^{14}t^7(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^2t^2+q^4t^4+q^6t^6+q^8t^8)\prod_{i=3}^{6}(1+q^{2i+6}t^{2i+1})+$$
$$q^{18}t^{10}(1-q^2(1+q^4t)(1+q^6t^3)(1+q^4t^3)(1+q^2t^2+q^4t^4+q^6t^6)(1+q^{14}t^9)(1+q^{16}t^{11})(1+q^{18}t^{13})+$$
$$q^{22}t^{14}(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^6t^3)(1+q^8t^5)(1+q^{16}t^{11})(1+q^{18}t^{13})(1+q^2t^2+2q^4t^4+q^6t^6+q^8t^8)+$$
$$q^{36}t^{25}(1-q^2)(1-q^4t^2)(1-q^6t^4)(1+q^4t)(1+q^6t^3)(1+q^8t^5)(1+q^{18}t^{13})(1+q^2t^2+q^4t^4)-$$
$$q^{42}t^{30}(1-q^2)(1-q^4t^2)(1+q^4t)(1+q^6t^3)(1+q^8t^5)(1+q^{10}t^7)(1+q^6t^5);$$

B Reduced Poincaré series

Here we collect the answers for the conjectural Poincaré series of stable reduced Khovanov homology for $n \leq 7$. The resulting series coincide with the data from [37] up to high q-degree. For example, for $(5,49)$ torus knot the first difference is in q-degree 100, for $(6,25)$ the first difference is in q-degree 52, for $(7,15)$ the first difference is in q-degree 32.

$$\overline{P}_3(q,t)=\frac{1+q^6t^3}{\prod_{i=1}^{2}(1-q^{2i+2}t^{2i})}[1-q^8t^4];$$

$$\overline{P}_4(q,t)=\frac{1+q^6t^3}{\prod_{i=1}^{3}(1-q^{2i+2}t^{2i})}[(1+q^{14}t^9)-q^8t^4(1+q^{14}t^9)-q^{10}t^6(1-q^4t^2)(1+q^8t^5)];$$

$$\overline{P}_5(q,t)=\frac{1+q^6t^3}{\prod_{i=1}^{4}(1-q^{2i+2}t^{2i})}[(1+q^{14}t^9)(1+q^{16}t^{11})-q^8t^4(1+q^{14}t^9)(1+q^{16}t^{11})-$$
$$q^{10}t^6(1-q^4t^2)(1+q^8t^5)(1+q^2t^2)(1+q^{16}t^{11})-q^{22}t^{15}(1-q^4t^2)(1-q^6t^4)(1+q^8t^5)];$$

$$\overline{P}_6(q,t)=\frac{1+q^6t^3}{\prod_{i=1}^{5}(1-q^{2i+2}t^{2i})}[(1+q^{14}t^9)(1+q^{16}t^{11})(1+q^{18}t^{13})-$$

$$q^8t^4(1+q^{14}t^9)(1+q^{16}t^{11})(1+q^{18}t^{13})-$$
$$q^{10}t^6(1-q^4t^2)(1+q^8t^5)(1+q^{16}t^{11})(1+q^{18}t^{13})(1+q^2t^2+q^4t^4)-$$
$$q^{22}t^{15}(1-q^4t^2)(1-q^6t^4)(1+q^8t^5)(1+q^{18}t^{13})(1+q^2t^2)+q^{26}t^{18}(1-q^4t^2)(1+q^8t^5)(1+q^{10}t^7)(1+q^6t^5)];$$

$$\overline{P}_7(q,t)=\frac{1+q^6t^3}{\prod_{i=1}^{6}(1-q^{2i+2}t^{2i})}[\prod_{i=1}^{4}(1+q^{2i+12}t^{2i+7})-q^8t^4\prod_{i=1}^{4}(1+q^{2i+12}t^{2i+7})-$$
$$q^{10}t^6(1-q^4t^2)(1+q^8t^5)(1+q^2t^2+q^4t^4+q^6t^6)\prod_{i=2}^{4}(1+q^{2i+12}t^{2i+7})-$$
$$q^{22}t^{15}(1-q^4t^2)(1-q^6t^4)(1+q^8t^5)(1+q^{18}t^{13})(1+q^{20}t^{15})(1+q^2t^2+q^4t^4)+$$
$$q^{26}t^{18}(1-q^4t^2)(1+q^8t^5)(1+q^{10}t^7)(1+q^2t^2)(1+q^{20}t^{15})(1+q^6t^5)+$$
$$q^{30}t^{22}(1-q^4t^2)(1-q^6t^4)(1+q^8t^5)(1+q^{10}t^7)(1+q^{12}t^9)].$$

C $(7,9)$ torus knot

We present the exact normalized Poincaré polynomial for the unreduced $\mathbb{Q}$-Khovanov homology of the $(7,9)$ torus knot, obtained with JavaKh ([6]).

$$q^{-47}P(T(7,8))=1+q^2+q^4t^2+q^8t^3+q^6t^4+q^8t^4+q^{10}t^5+q^{12}t^5+q^8t^6+q^{10}t^6+q^{12}t^7+q^{14}t^7+q^{10}t^8+$$
$$2q^{12}t^8+q^{14}t^9+2q^{16}t^9+q^{12}t^{10}+2q^{14}t^{10}+q^{16}t^{11}+3q^{18}t^{11}+q^{14}t^{12}+3q^{16}t^{12}+q^{18}t^{12}+$$
$$q^{22}t^{12}+q^{18}t^{13}+4q^{20}t^{13}+q^{22}t^{13}+3q^{18}t^{14}+q^{20}t^{14}+q^{24}t^{14}+5q^{22}t^{15}+2q^{24}t^{15}+2q^{20}t^{16}+$$
$$3q^{22}t^{16}+2q^{26}t^{16}+q^{28}t^{16}+4q^{24}t^{17}+4q^{26}t^{17}+q^{22}t^{18}+3q^{24}t^{18}+2q^{28}t^{18}+q^{30}t^{18}+2q^{26}t^{19}+$$
$$5q^{28}t^{19}+q^{24}t^{20}+3q^{26}t^{20}+q^{30}t^{20}+2q^{32}t^{20}+q^{28}t^{21}+5q^{30}t^{21}+2q^{28}t^{22}+q^{30}t^{22}+2q^{34}t^{22}+$$
$$5q^{32}t^{23}+q^{34}t^{23}+q^{30}t^{24}+3q^{32}t^{24}+3q^{36}t^{24}+2q^{34}t^{25}+4q^{36}t^{25}+q^{34}t^{26}+q^{36}t^{26}+2q^{38}t^{26}+$$
$$q^{40}t^{26}+3q^{38}t^{27}+q^{40}t^{27}+q^{42}t^{27}+q^{38}t^{28}+2q^{42}t^{28}+2q^{42}t^{29}+q^{42}t^{30}+q^{46}t^{30}+q^{46}t^{31}.$$

The total dimension of the homology equals to 134. One can observe the multiplicative generators of the following (bi)degrees:

$$\deg(x_0)=q^2,\ \deg(x_1)=q^4t^2,\ \deg(x_2)=q^6t^4,\ \deg(x_3)=q^8t^6,$$
$$\deg(x_4)=q^{10}t^8,\ \deg(x_5)=q^{12}t^{10},\ \deg(x_6)=q^{14}t^{12},$$
$$\deg(\mu_0)=q^8t^3,\ \deg(\mu_1)=q^{10}t^5,\ \deg(\mu_2)=q^{12}t^7,$$
$$\deg(\mu_3)=q^{14}t^9,\ \deg(\mu_4)=q^{16}t^{11},\ \deg(\mu_5)=q^{18}t^{13}.$$

Let us consider $\mathbb{Z}_2$ coefficients. The Poincaré polynomial equals to:

$$
\begin{gathered}
q^{-47}P_2(T(7,9)) = 1+q^2+q^4t^2+q^6t^2+q^6t^3+q^8t^3+q^6t^4+q^8t^4+q^{10}t^5+q^{12}t^5+q^8t^6+2q^{10}t^6+\\
q^{12}t^6+q^{10}t^7+2q^{12}t^7+q^{14}t^7+q^{10}t^8+2q^{12}t^8+q^{14}t^8+2q^{14}t^9+3q^{16}t^9+q^{18}t^9+q^{12}t^{10}+3q^{14}t^{10}+\\
3q^{16}t^{10}+q^{18}t^{10}+q^{14}t^{11}+3q^{16}t^{11}+3q^{18}t^{11}+q^{20}t^{11}+q^{14}t^{12}+3q^{16}t^{12}+3q^{18}t^{12}+2q^{20}t^{12}+\\
q^{22}t^{12}+3q^{18}t^{13}+6q^{20}t^{13}+3q^{22}t^{13}+3q^{18}t^{14}+5q^{20}t^{14}+3q^{22}t^{14}+q^{24}t^{14}+3q^{20}t^{15}+6q^{22}t^{15}+\\
4q^{24}t^{15}+q^{26}t^{15}+2q^{20}t^{16}+4q^{22}t^{16}+4q^{24}t^{16}+3q^{26}t^{16}+q^{28}t^{16}+q^{22}t^{17}+7q^{24}t^{17}+7q^{26}t^{17}+\\
q^{28}t^{17}+q^{22}t^{18}+5q^{24}t^{18}+6q^{26}t^{18}+3q^{28}t^{18}+q^{30}t^{18}+q^{24}t^{19}+5q^{26}t^{19}+7q^{28}t^{19}+3q^{30}t^{19}+\\
q^{24}t^{20}+3q^{26}t^{20}+5q^{28}t^{20}+6q^{30}t^{20}+3q^{32}t^{20}+4q^{28}t^{21}+9q^{30}t^{21}+5q^{32}t^{21}+2q^{28}t^{22}+\\
5q^{30}t^{22}+5q^{32}t^{22}+3q^{34}t^{22}+q^{36}t^{22}+2q^{30}t^{23}+6q^{32}t^{23}+6q^{34}t^{23}+2q^{36}t^{23}+q^{30}t^{24}+3q^{32}t^{24}+\\
6q^{34}t^{24}+5q^{36}t^{24}+q^{38}t^{24}+4q^{34}t^{25}+6q^{36}t^{25}+2q^{38}t^{25}+q^{34}t^{26}+3q^{36}t^{26}+4q^{38}t^{26}+2q^{40}t^{26}+\\
q^{36}t^{27}+4q^{38}t^{27}+4q^{40}t^{27}+q^{42}t^{27}+q^{38}t^{28}+3q^{40}t^{28}+2q^{42}t^{28}+q^{40}t^{29}+2q^{42}t^{29}+q^{44}t^{29}+\\
q^{42}t^{30}+2q^{44}t^{30}+q^{46}t^{30}+q^{44}t^{31}+q^{46}t^{31}.
\end{gathered}
$$

The total dimension of the homology is 286 (about twice bigger than for $\mathbb{Q}$-coefficients), and the bidegrees of the multiplicative generators are equal to:

$$
\deg(x_0) = q^2,\ \deg(x_1) = q^4t^2,\ \deg(x_2) = q^6t^4,\ \deg(x_3) = q^8t^6,
$$

$$
\begin{gathered}
\deg(x_4) = q^{10}t^8,\ \deg(x_5) = q^{12}t^{10},\ \deg(x_6) = q^{14}t^{12},\\
\deg(\xi_1) = q^6t^3,\ \deg(\xi_3) = q^{10}t^7,\ \deg(\xi_5) = q^{14}t^1 1.
\end{gathered}
$$

Finally, one can check that the Khovanov homology of this knot has nontrivial $\mathbb{Z}_7$-torsion in degree $q^{20}t^{14}$.

References

[1] M. Asaeda, J. Przytycki. Khovanov homology: torsion and thickness. Proceedings of the Workshop, "New Techniques in Topological Quantum Field Theory" Calgary/Kananaskis, Canada, August 2001; Ed. J.Bryden.

[2] G. E. Andrews. The theory of partitions. Addison-Wesley, 1976.

[3] D. Bar-Natan. On Khovanov's categorification of the Jones polynomial. Algebraic and Geometric Topology **2** (2002) 337-370.

[4] D. Bar-Natan. Khovanov's homology for tangles and cobordisms. Geom. Topol. **9**(2005) 1443-99.

[5] D. Bar-Natan. Fast Khovanov Homology Computations. J. Knot Theory Ramifications **16** (2007), no. 3, 243–255.

[6] D. Bar-Natan, S. Morrisson. The Knot Atlas. `http://katlas.org`

[7] C. Bruschek, H. Mourtada, J. Schepers. Arc Spaces and Rogers-Ramanujan Identities. 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), 211220, Discrete Math. Theor. Comput. Sci. Proc., AO, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2011.

[8] W. Decker, G.-M. Greuel, G. Pfister, H. Schönemann. SINGULAR 3-1-3 — A computer algebra system for polynomial computations. http://www.singular.uni-kl.de (2011).

[9] N. Dunfield, S. Gukov, J. Rasmussen.The Superpolynomial for Knot Homologies. Experimental Math. **15** (2006), 129–159.

[10] T. Dyckerhoff. Compact generators in categories of matrix factorizations. Duke Math. J. **159** (2011), no. 2, 223–274.

[11] D. Eisenbud. Homological algebra on a complete intersection, with an application to group representations. Trans. Amer. Math. Soc. **260**, 35–64, 1980.

[12] B. Feigin. Abelianization of the BGG resolution of representations of the Virasoro algebra. Funct. Anal. Appl. **45** (2011), no. 4, 297–304.

[13] B. Feigin, A. Stoyanovsky. Functional models for representations of current algebras and semi-infinite Schubert cells. Funct. Anal. Appl. **28** (1994), no. 1, 55-72.

[14] B. Feigin, S. Loktev. On the finitization of Gordon identities. Funct. Anal. Appl. **35** (2001), no. 1, 44–51.

[15] I. Frenkel, C. Stroppel, J. Sussan. Categorifying fractional Euler characteristics, Jones-Wenzl projector and $3j$-symbols. arXiv:1007.4680

[16] E. Gorsky. q, t–Catalan numbers and knot homology. Zeta Functions in Algebra and Geometry, 213–232. Contemp. Math. **566**, Amer. Math. Soc., Providence, RI, 2012.

[17] E. Gorsky, A. Oblomkov, J. Rasmussen, V. Shende. Khovanov-Rozansky homology of torus knots and simple representations of rational Cherednik algebras , in preparation.

[18] S. Gukov, J. Walcher. Matrix Factorizations and Kauffman Homology. arXiv: 0512298

[19] S. Gukov, M. Stošić. Homological algebra of knots and BPS states. arXiv:1112.0030

[20] G. Hochschild, B. Kostant, A. Rosenberg, Differential forms on regular affine algebras, Transactions AMS **102** (1962), No.3, 383–408.

[21] V. Jones. Hecke algebra representations of braid groups and link polynomials. Ann. of Math. **126** (1987), no.2, 335–388.

[22] V. Kac. Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990.

[23] M. Khovanov. A categorification of the Jones polynomial. Duke Math. J. **101** (2000), no. 3, 359–426.

[24] M. Khovanov. Patterns in knot cohomology I. Experimental Math. **12** (2003) no.3, 365–374.

[25] M. Khovanov. Triply-graded link homology and Hochschild homology of Soergel bimodules. Int. Journal of Math. **18** (2007), no. 8 869–885.

[26] M. Khovanov, L. Rozansky. Matrix factorizations and link homology I. Fund. Math. **199** (2008), no. 1, 1–91.

[27] M. Khovanov, L. Rozansky. Matrix factorizations and link homology II. Geom. Topol. **12** (2008), no. 3, 1387–1425.

[28] E. S. Lee. An endomorphism of the Khovanov invariant. Adv. Math. **197** (2005), no. 2, 554–586.

[29] J. Lepowsky, M. Primc. Structure of the standard modules for the affine Lie algebra $A_1^{[1]}$. Contemporary Mathematics, 46. American Mathematical Society, Providence, RI, 1985.

[30] S. Loktev. Fermionic formulas for the characters of the representations of $\widehat{\mathrm{sl}}_2$. Russian Math. Surveys **56** (2001), no. 1, 164–165.

[31] A. Oblomkov, V. Shende. The Hilbert scheme of a plane curve singularity and the HOMFLY polynomial of its link. Duke Math. J. **161** (2012), no. 7 , 1277–1303.

[32] A. Oblomkov, J. Rasmussen, V. Shende. The Hilbert scheme of a plane curve singularity and the HOMFLY homology of its link. arXiv:1201.2115

[33] J. Rasmussen. Some differentials on Khovanov-Rozansky homology. arXiv: math.GT/0607544

[34] J. Rasmussen. Khovanov homology and the slice genus. Invent. Math. **182** (2010), no. 2, 419–447.

[35] L. Rozansky. An infinite torus braid yields a categorified Jones-Wenzl projector. arXiv:1005.3266

[36] A. Shumakovitch. KhoHo: a program for computing Khovanov homology. `www.geometrie.ch/KhoHo` (currently unavailable).

[37] A. Shumakovitch. Private communication.

[38] A. Shumakovitch. Torsion of the Khovanov homology. arXiv:math/0405474

[39] M. Stošić. Khovanov homology of torus links. Topology Appl. **156** (2009), no. 3, 533–541

[40] M. Stošić. Homological thickness and stability of torus knots. Algebr. Geom. Topol. **7** (2007), 261–284.

[41] C. Stroppel, J. Sussan. Categorified Jones-Wenzl Projectors: a comparison. arXiv:1105.3038

[42] P. Turner. A spectral sequence for Khovanov homology with an application to $(3, q)$-torus links. Algebr. Geom. Topol. **8** (2008), no. 2, 869–884.

[43] H. Wenzl. On sequences of projections. C. R. Math. Rep. Acad. Sci. Canada **9** (1987), no. 1, 5–9.

[44] B. Webster. Knot invariants and higher representation theory II: the categorification of quantum knot invariants. arXiv:1005.4559

Eugene Gorsky
Department of Mathematics
Stony Brook University
Stony Brook, NY 11794
`egorsky@math.sunysb.edu`

Alexei Oblomkov
Department of Mathematics
University of Massachusetts, Amherst
Amherst, MA 01003
`oblomkov@math.umass.edu`

Jacob Rasmussen
Department of Pure Mathematics
University of Cambridge
Centre for Mathematics Sciences
Wilberforce Road, CB3 0WB
United Kingdom
`J.Rasmussen@dpmms.cam.ac.uk`